

On Minimum Sets of 1-Factors Covering a Complete Multipartite Graph

David Cariolaro¹ and Hung-Lin Fu²

¹INSTITUTE OF MATHEMATICS
ACADEMIA SINICA, NANKANG, TAIPEI
11529, TAIWAN

E-mail: cariolaro@math.sinica.edu.tw; davidcariolaro@hotmail.com

²DEPARTMENT OF APPLIED MATHEMATICS
NATIONAL CHIAO TUNG UNIVERSITY
HSIN CHU 30050
TAIWAN

E-mail: hlfu@math.nctu.edu.tw

Received April 20, 2006; Revised January 21, 2008

Published online 15 April 2008 in Wiley InterScience(www.interscience.wiley.com).
DOI 10.1002/jgt.20303

Abstract: We determine necessary and sufficient conditions for a complete multipartite graph to admit a set of 1-factors whose union is the whole graph and, when these conditions are satisfied, we determine the minimum size of such a set. © 2008 Wiley Periodicals, Inc. *J Graph Theory* 58: 239–250, 2008

Keywords: *1-factor; 1-factor cover; excessive factorization; excessive index; complete multipartite graph*

1. INTRODUCTION

All graphs considered will be finite, simple and undirected, unless stated otherwise. We denote by $V(G)$ and $E(G)$, respectively, the vertex and edge set of a graph G .

Journal of Graph Theory
© 2008 Wiley Periodicals, Inc.



The *order* of G is $|V(G)|$. The *maximum degree* of G will be denoted by $\Delta(G)$ and its *chromatic index* by $\chi'(G)$.

Let G be a graph of even order. A *1-factor* (or *perfect matching*) of G is a 1-regular spanning subgraph, that is, a set of exactly $\frac{|V(G)|}{2}$ independent edges.¹ G is *1-extendable* if every edge of G belongs to at least one 1-factor of G . A *1-factor cover* of G is a set \mathcal{F} of 1-factors of G such that $\cup_{F \in \mathcal{F}} = E(G)$. Notice that G admits a 1-factor cover if and only if it is 1-extendable. If G is 1-extendable, a 1-factor cover of minimum cardinality will be called an *excessive factorization*.

Thus, a *1-factorization* of G is a 1-factor cover \mathcal{F} with the property that all the 1-factors in \mathcal{F} are pairwise disjoint. Any 1-factorization is an excessive factorization, but the converse is obviously not true. For example, the Petersen graph has no 1-factorization, but has an excessive factorization consisting of five 1-factors (see [2]). The graphs which admit an excessive factorization are precisely those that have a 1-factor cover, that is, those that are 1-extendable.

Let G be a 1-extendable graph. The *excessive index* of G , denoted $\chi'_e(G)$, is the size of an excessive factorization of G . We define $\chi'_e(G) = \infty$ if G is not 1-extendable.

Bonisoli [1] and Wallis [6] considered 1-factor covers of the complete graph K_{2n} which do not contain a 1-factorization of K_{2n} .

Bonisoli and Cariolaro [2] introduced the concept of *excessive factorization*, defined the parameter $\chi'_e(G)$, and studied excessive factorizations of regular graphs. They posed a number of open problems and conjectures. A first question is, of course, to determine $\chi'_e(G)$ for any graph G . It is observed in [2] that this problem is NP-hard since, if G is regular and has even order, then $\chi'_e(G) = \Delta(G)$ if and only if G is 1-factorizable, and to determine whether a regular graph G is 1-factorizable is NP-complete. Therefore, we can expect to be able to determine $\chi'_e(G)$ only for some specific classes of graphs.

In this article, we consider the class of *complete multipartite* graphs. Hoffman and Rodger [3] determined the chromatic index of all complete multipartite graphs. Here, we shall determine the excessive index $\chi'_e(G)$ of any complete multipartite graph G .

We will often use, without further reference, the following fact, proved by de Werra [7] and, independently, by McDiarmid [4]. If a multigraph G has a k -edge coloring, that is, if $k \geq \chi'(G)$, then it also has an *equalized* k -edge coloring, namely a k -edge coloring such that each color class has size either $\lfloor \frac{|E(G)|}{k} \rfloor$ or $\lceil \frac{|E(G)|}{k} \rceil$.

We shall also need the concept of *excessive coloring*. An excessive coloring of a graph G is an assignment of (possibly more than one) colors to each of the edges of G such that the edges on which a given color appears are independent (i.e., they form a matching). Thus an excessive coloring can be simply specified

¹To be precise, a 1-factor F of G is a 1-regular spanning subgraph of G and a perfect matching is the edge set of F , but the two terms are often used interchangeably in the literature.

as a collection of matchings of G whose union is $E(G)$. It is normally interesting to consider this concept when additional restrictions are imposed on the matchings which form the color classes. For example, when each color class is a 1-factor, the corresponding excessive coloring is equivalent to a 1-factor cover.

2. 1-EXTENDABLE COMPLETE MULTIPARTITE GRAPHS

We adopt the notation $G = K(n_1, n_2, \dots, n_r)$ to designate a complete multipartite graph with partite sets of size n_1, n_2, \dots, n_r , where $n_1 \geq n_2 \geq n_3 \geq \dots \geq n_r$. We also let V_1, V_2, \dots, V_r denote the r partite sets of G . By definition, for each i , V_i is an independent set of n_i vertices of G which are joined to every vertex in $G - V_i$.

Trivially, the complete bipartite graph $K(m, n)$ is 1-extendable if and only if $m = n$, in which case it actually has a 1-factorization. Therefore, $\chi'_e(K(m, n)) = n$ if $n = m$ and ∞ otherwise. From now on we make the convention that all complete multipartite graphs considered have r partite sets, where $r \geq 3$. The following lemma is probably well known, but we give a full proof for the sake of completeness.

Lemma 1. *The graph $G = K(n_1, n_2, \dots, n_r)$ has a 1-factor if and only if*

1. $\sum_{i=1}^r n_i$ is even;
2. $n_1 \leq \sum_{i=2}^r n_i$.

Proof. The first of the above conditions is clearly necessary in order for the graph G to have a 1-factor, as G must have even order. To see the necessity of the second, it suffices to see that any 1-factor of G must match the vertices of V_1 (the first partite set) to the vertices of the complement (since the vertices in V_1 are mutually nonadjacent). Hence, the two conditions are necessary. To see the sufficiency, assume both conditions hold. We prove the existence of a 1-factor by induction on k , where $2k = \sum_{i=2}^r n_i - n_1$.

If $k = 0$, then we have $n_1 = \sum_{i=2}^r n_i$ and a 1-factor of G is easily obtained by matching the vertices of V_1 to the vertices of $V_2 \cup V_3 \cup \dots \cup V_r$. Assume now that the theorem holds for any G with $\sum_{i=2}^r n_i - n_1 < 2k$ and consider the case of a G with $\sum_{i=2}^r n_i - n_1 = 2k$. Let $x \in V_r$ and $y \in V_{r-1}$ and consider the edge $e = xy$. We prove that there is a 1-factor of G containing this edge. This is equivalent to proving that the graph $G - x - y$ has a 1-factor. But it is easily seen that the graph $G - x - y$ is complete multipartite with partite sets V'_1, V'_2, \dots, V'_r , where $|V'_i| = n_i$ for all $i \leq r - 2$ and $|V'_{r-1}| = n_{r-1} - 1$ and $|V'_r| = n_r - 1$. Moreover, $G - x - y$ satisfies the inductive hypothesis, since $|V'_2| + |V'_3| + \dots + |V'_{r-1}| + |V'_r| - |V'_1| = 2k - 2$. Thus, $G - x - y$ has a 1-factor and hence G has the desired 1-factor. ■

Using Lemma 1, it is easy to determine which complete multipartite graphs admit an excessive factorization, as given by the following theorem.

Theorem 1. *The graph $K(n_1, n_2, \dots, n_r)$ is 1-extendable if and only if*

1. $\sum_{i=1}^r n_i$ is even;
2. $n_1 < \sum_{i=2}^r n_i$.

Proof. Let $G = K(n_1, n_2, \dots, n_r)$ and suppose G is 1-extendable. The first condition follows immediately from the fact that G has a 1-factor. Let e be an edge joining the second and third partite sets. By the fact that G is 1-extendable, there exists a 1-factor F containing e . Clearly, this 1-factor must match the vertices of the first partite set onto the vertices of the complement, but the two vertices which are the endpoint of e are F -saturated. Hence, the condition (2) above (which is clearly equivalent to $n_1 \leq \sum_{i=2}^r n_i - 2$, given the parity of G) must hold.

Conversely, suppose G satisfies both conditions above. We prove that G is 1-extendable. Let $e \in E(G)$. We prove the existence of a 1-factor F containing e . Equivalently, we prove that the graph $G - x - y$ has a 1-factor, where $xy = e$. Without loss of generality, we can assume $x \in V_i, y \in V_j$, and $i < j$. But then $G - x - y \cong G_1 = K(n_1, n_2, \dots, n_{i-1}, n_i - 1, n_{i+1}, \dots, n_{j-1}, n_j - 1, n_{j+1}, \dots, n_r)$ and G_1 is easily seen to satisfy the hypotheses of Lemma 1. Hence, G_1 has a 1-factor and G has the desired 1-factor. ■

3. SOME LEMMAS

Theorem 1 gives necessary and sufficient conditions for a complete multipartite graph G to admit an excessive factorization, that is, to satisfy $\chi'_e(G) < \infty$. We are now left with the task of determining precisely $\chi'_e(G)$ for all such graphs.

We start by giving some lower bounds. One obvious lower bound is the maximum degree of G , because every edge incident with a vertex of maximum degree must belong to a distinct 1-factor in an excessive factorization. Thus, $\chi'_e(G) \geq \Delta(G)$ holds, not only for complete multipartite graphs, but for all graphs G (in fact an easy argument along the same line also shows that, for all graphs G , $\chi'_e(G) \geq \chi'(G)$, but we shall not need this stronger inequality here).

The next lower bound is less trivial. Let $G = K(n_1, n_2, \dots, n_r)$. Let V_i be the i th partite set of G . Let E_i be defined as

$$E_i = E(G - V_i).$$

Define

$$\sigma_i(G) = \left\lceil \frac{2|E_i|}{|V(G)| - 2|V_i|} \right\rceil. \quad (1)$$

Since the vertices of V_i are independent in G , any 1-factor F of G must contain exactly n_i edges joining V_i to $G - V_i$. Thus, in particular, the 1-factor F contains exactly $\frac{|V(G)| - 2|V_i|}{2}$ edges from E_i . It follows from (1) that any 1-factor cover must

contain at least $\sigma_i(G)$ 1-factors. This proves that

$$\chi'_e(G) \geq \max_{1 \leq i \leq r} \sigma_i(G).$$

As shown next, the quantity $\max_{1 \leq i \leq r} \sigma_i(G)$ is particularly simple to evaluate, since it is always equal to $\sigma_1(G)$.

Proposition 1. *Let $G = K(n_1, n_2, \dots, n_r)$. Then*

$$\sigma_1(G) = \max_{1 \leq i \leq r} \sigma_i(G),$$

where the parameters $\sigma_i(G)$ are defined in (1).

Proof. We start by observing that

$$\sigma_k(G) = \frac{2 \sum_{1 \leq i < j \leq r, i, j \neq k} n_i n_j}{\sum_{i=1}^r n_i - 2n_k}.$$

Therefore, Proposition 1 follows from the truth of the following inequality involving positive integers x_1, x_2, \dots, x_r , where $x_1 = \max_{1 \leq i \leq r} x_i$:

$$\frac{\sum_{2 \leq i < j \leq r} x_i x_j}{\sum_{i=1}^r x_i - 2x_1} \geq \frac{\sum_{1 \leq i < j \leq r, i, j \neq k} x_i x_j}{\sum_{i=1}^r x_i - 2x_k}. \tag{2}$$

By the arbitrariness of the x_i s ($i > 1$), we can assume $k = 2$. Further, if $x_1 = x_2$ there is clearly nothing to prove because the two sides of (2) are in this case identical. Thus, we may assume that $x_1 > x_2$. We have to prove that

$$\left(\sum_{2 \leq i < j \leq r} x_i x_j \right) \left(\sum_{i=1}^r x_i - 2x_2 \right) \geq \left(\sum_{1 \leq i < j \leq r, i, j \neq 2} x_i x_j \right) \left(\sum_{i=1}^r x_i - 2x_1 \right).$$

Let $\xi = x_3 + x_4 + \dots + x_r$ and let $\eta = \sum_{3 \leq i < j \leq r} x_i x_j$. Using these notations, we can rewrite the above inequality as

$$(x_2 \xi + \eta)(x_1 - x_2 + \xi) \geq (x_1 \xi + \eta)(x_2 - x_1 + \xi).$$

Multiplying out, simplifying and rearranging the terms, we obtain

$$(x_1 - x_2)\xi^2 - (x_1^2 - x_2^2)\xi - 2\eta(x_1 - x_2) \leq 0.$$

Dividing by $x_1 - x_2$, which is positive by assumption, we obtain

$$\xi^2 - (x_1 + x_2)\xi - 2\eta \leq 0.$$

But we have

$$\begin{aligned} \xi^2 &= (x_3 + x_4 + \dots + x_r)^2 = x_3^2 + x_4^2 + \dots + x_r^2 + 2 \\ &\quad \cdot \sum_{3 \leq i < j \leq r} x_i x_j = x_3^2 + x_4^2 + \dots + x_r^2 + 2\eta. \end{aligned}$$

Thus, the problem is reduced to showing that

$$x_3^2 + x_4^2 + \cdots + x_r^2 \leq (x_1 + x_2)(x_3 + x_4 + \cdots + x_r).$$

Using the assumption that $x_1 = \max_{1 \leq i \leq r} x_i$ and the fact that

$$x_3^2 + x_4^2 + \cdots + x_r^2 \leq x_1(x_3 + x_4 + \cdots + x_r) < (x_1 + x_2)(x_3 + x_4 + \cdots + x_r),$$

we conclude the proof. ■

Thus, we have two nontrivial lower bounds on $\chi'_e(G)$, one is $\Delta(G)$ and the other is $\sigma_1(G)$. Neither of the two is necessarily worse or better than the other. For example, if $G = K(5, 4, 3)$ then $\sigma_1(G) = 12 > \Delta(G) = 9$, but if $G = K(4, 4, 4, 2)$ then $\sigma_1(G) = 11 < \Delta(G) = 12$.

Let

$$\tau(G) = \max\{\sigma_1(G), \Delta(G)\}.$$

By what we have just proved, we have

Lemma 2. *Let G be a 1-extendable complete r -partite graph. Then*

$$\chi'_e(G) \geq \tau(G).$$

In the next section, we shall prove that the above inequality is indeed an equality by proving the following theorem.

Theorem 2. *Let G be a 1-extendable complete r -partite graph. Then*

$$\chi'_e(G) = \tau(G) = \max\{\sigma_1(G), \Delta(G)\}.$$

We shall use the following lemma, which can be seen as an extension of Hall's Theorem. It generalizes a theorem of Bondy ([5, Theorem 13.3, p.109]). *Inter alia*, it completely solves the problem of characterizing the excessive colorings of $G - V_1$ which extend to 1-factor covers of G , for any complete multipartite graph G .

Lemma 3. *Let \mathcal{C} be a set (of colors) and let s, t be positive integers, with $s \leq t \leq |\mathcal{C}|$. Let $T = \{y_1, y_2, \dots, y_t\}$ be a set of cardinality t and let $S = \{x_1, x_2, \dots, x_s\}$ be a set (disjoint from T) of cardinality s . For each $y \in T$, let $L(y) \subset \mathcal{C}$ be a set of colors. Let, for each $\alpha \in \mathcal{C}$, $T_\alpha = \{y \in T; \alpha \in L(y)\}$. Consider the complete bipartite graph X with bipartition (T, S) . There exists an excessive coloring ψ of X with color set \mathcal{C} such that, for each $\alpha \in \mathcal{C}$, the color class corresponding to α is a perfect matching from S to T_α if and only if the following conditions are satisfied:*

1. every $\alpha \in \mathcal{C}$ is contained in precisely s sets of the family $\{L(y) \mid y \in T\}$;
2. $|L(y)| \geq s$ (for all $y \in T$).

Proof. Assume that there exists an excessive coloring as in the statement of the lemma. Then clearly condition (1) is satisfied since the existence of a perfect matching from S to T_α implies $|T_\alpha| = s$. Condition (2) follows from the fact that

every $y \in T$ is incident with s edges in the graph X , and each of these edges must be assigned at least one distinct color by ψ , this color being in $L(y)$. Thus, the two conditions above are necessary for the existence of ψ . We now show that they are sufficient.

Consider the bipartite graph B_1 with bipartition (T, C) , where there is an edge between y and α if and only if $\alpha \in L(y)$. By assumption, $\deg_{B_1}(\alpha) = s$ for all $\alpha \in C$ and $\deg_{B_1}(y) \geq s$ for all $y \in T$.

Let B_2 be a spanning subgraph of B_1 such that

$$\deg_{B_2}(y) = s \text{ for all } y \in T.$$

Since B_2 is bipartite and $\Delta(B_2) = s$, by König's Theorem B_2 has an s -edge coloring π with colors $\{1, 2, \dots, s\}$. We now define an edge coloring θ of X as follows: if y is joined in B_2 to color α by an edge colored j , we color the edge yx_j of X by color α . It is easy to see that the coloring θ is well defined, and that it is in fact a proper edge coloring of X . To obtain the required excessive coloring of X , it is now sufficient, for each color α , to extend arbitrarily the color class C_α corresponding to α to a perfect matching from S to T_α . ■

Instead of proving Theorem 2 directly, in the next section we shall prove the following theorem, whose equivalence with Theorem 2 will be established below.

Theorem 3. *Let $G = K(n_1, n_2, \dots, n_r)$ be a 1-extendable complete r -partite graph and let $H = K(n_2, n_3, \dots, n_r)$. Then there exists an excessive coloring ϕ of H with exactly $\tau(G)$ colors such that each color class misses exactly n_1 vertices of H and each vertex of H misses at least n_1 colors.*

Using Lemma 3, we now prove the equivalence between Theorems 2 and 3.

Lemma 4. *Theorem 2 holds for G if and only if Theorem 3 does.*

Proof. Assume Theorem 2 is true for the graph $G = K(n_1, n_2, \dots, n_r)$. Let ψ be an excessive factorization of G . Then ψ , when viewed as an excessive coloring, consists of $\tau(G)$ color classes. Let $H = G - V_1$, where V_1 is the largest partite set of G . Then $H \cong K(n_2, n_3, \dots, n_r)$. Clearly, the restriction ϕ of ψ to $E(H)$ makes Theorem 3 true for the graph G . Hence, if Theorem 2 holds for G then Theorem 3 does. For the converse, let G and H be as above and assume Theorem 3 holds for G . By Theorem 3, there exists an excessive coloring ϕ of H with $\tau(G)$ color classes such that each color class is a matching missing exactly n_1 vertices of H and each vertex of H misses at least n_1 colors. We now extend ϕ to an excessive coloring of G as follows. Let \mathcal{C} be the color set of ϕ . For each vertex $v \in V(H)$, let $L(v)$ be the set of colors missing at v , that is, the set of colors which do not appear on any of the edges incident with v . By the conditions satisfied by ϕ , $|L(v)| \geq n_1$ for all $v \in V(H)$ and every color $\alpha \in \mathcal{C}$ appears on exactly n_1 of the sets $L(v)$, $v \in V(H)$. Therefore, the conditions of Lemma 3 are satisfied by the color set \mathcal{C} , the family of sets $\mathcal{L} = \{L(v) \mid v \in V(H)\}$ and the set $S = V_1$. By Lemma 3, there exists an excessive coloring ψ of the complete bipartite graph U with bipartition $(V(H), V_1)$

such that, for every color α , the color class corresponding to α is a perfect matching from S to T_α , where T_α is the set of vertices in $V(H)$ which are missing (with respect to ϕ) color α . Therefore, it is easily seen that the map ρ defined by

$$\rho(e) = \begin{cases} \phi(e) & \text{if } e \in E(H), \text{ and} \\ \psi(e) & \text{if } e \in E(U). \end{cases}$$

is an excessive coloring of G which uses $\tau(G)$ colors and such that every color class is a 1-factor of G . Therefore, ρ is a 1-factor cover of G consisting of $\tau(G)$ 1-factors, and, by Lemma 2, this number is necessarily the minimum, which proves that ρ is an excessive factorization of G . Thus, Theorem 2 holds for G . This concludes the proof of Lemma 4. ■

4. THE MAIN RESULT

We now prove Theorem 2 for all those complete multipartite graphs for which $\sigma_1(G) > \Delta(G)$ by proving the following.

Theorem 4. *Let G be a 1-extendable complete multipartite graph such that $\sigma_1(G) > \Delta(G)$. Then $\chi'_e(G) = \sigma_1(G)$.*

Proof. Let $G = K(n_1, n_2, \dots, n_r)$ and let $H = G - V_1 \cong K(n_2, n_3, \dots, n_r)$, where V_1 is the largest partite set of G , and assume that G is 1-extendable and $\sigma_1(G) > \Delta(G)$. By Lemma 4, it will suffice to show that Theorem 3 holds for G . Thus, we need to find an excessive coloring ϕ of H with exactly $\sigma_1(G)$ color classes, each of which misses exactly n_1 vertices of H and with respect to which each vertex of H misses at least n_1 colors. Let $m = \frac{|V(H)| - n_1}{2}$. Notice that $\sigma_1(G) = \lceil \frac{|E(H)|}{m} \rceil$. Let $m_1 = |E(H)| - (\sigma_1(G) - 1)m$. Notice that $0 < m_1 \leq m$.

We prove that there exists an edge-coloring of H with $\sigma_1(G) - 1$ color classes of size m and 1 color class of size m_1 .

By assumption G has a 1-factor, and hence H has a matching of size m . Let M_1 be a matching in H of size m_1 . Consider the graph $H - M_1$. We have

$$\chi'(H - M_1) \leq \chi'(H) \leq \chi'(G) = \Delta(G) \leq \sigma_1(G) - 1.$$

Hence, there exists a $(\sigma_1(G) - 1)$ -edge coloring of $H - M_1$. Notice that $|E(H - M_1)| = (\sigma_1(G) - 1)m$. But then there exists an *equalized* $(\sigma_1(G) - 1)$ -edge coloring of $H - M_1$, so that each color class has size exactly m .

Putting back the matching M_1 as an additional color class, we have the required edge coloring of H . But now, in order to obtain an excessive coloring of H as in Theorem 3, we just need to extend the color class M_1 to an arbitrary color class (matching) of size m of H . The excessive coloring ϕ thus defined is such that any vertex v of H of degree $\deg_H(v)$ “sees” at most $\deg_H(v) + 1$ colors, because the only possible edges with multiple colors are those in M_1 . Thus, at any vertex of H there are at most

$$\sigma_1(G) - (\deg_H(v) + 1) \geq \sigma_1(G) - \Delta(H) - 1 \geq \Delta(G) - \Delta(H) = n_1$$

colors missing. Notice that each color class of ϕ is a matching of size m and hence (by the definition of m) misses exactly n_1 vertices of H . Therefore, ϕ verifies Theorem 3 for G , and hence (by Lemma 4) it verifies Theorem 2 for G . This terminates the proof. ■

To terminate the proof of Theorem 2, we need to settle the case $\sigma_1(G) \leq \Delta(G)$. The following notation will be helpful in the sequel. If Z is a graph, $v \in V(Z)$ and k is an integer, the k -deficiency of v in Z is the quantity

$$k_def(v) = k - \deg_Z(v)$$

and the k -deficiency of Z is defined as

$$k_def(Z) = \sum_{v \in V(Z)} k_def(v) = \sum_{v \in V(Z)} (k - \deg_Z(v)).$$

Notice that, if $k = \Delta(G)$, then the k -deficiency of Z is usually called *deficiency* of Z and denoted by $def(Z)$.

Proposition 2. *Let $G = K(n_1, n_2, \dots, n_r)$ and let $H = G - V_1 \cong K(n_2, n_3, \dots, n_r)$. Then the following two conditions are equivalent:*

1. $\sigma_1(G) \leq \Delta(G)$;
2. $\Delta(G)_def(H) \geq n_1 \Delta(G)$.

Proof. Condition 1 is equivalent to

$$\frac{2|E(H)|}{|V(H)| - n_1} \leq \Delta(G),$$

that is,

$$2|E(H)| \leq (|V(H)| - n_1)\Delta(G),$$

that is,

$$\sum_{v \in V(H)} \deg_H(v) \leq (|V(H)| - n_1)\Delta(G),$$

that is,

$$\sum_{v \in V(H)} (\deg_H(v) - \Delta(G)) \leq -n_1 \Delta(G).$$

Changing sign, we have

$$\Delta(G)_def(H) \geq n_1 \Delta(G)$$

which is condition 2. ■

We will need the following lemma.

Lemma 5. *Let $G = K(n_1, n_2, \dots, n_r)$ be a 1-extendable complete multipartite graph such that $\sigma_1(G) \leq \Delta(G)$ and let $H = G - V_1 \cong K(n_2, n_3, \dots, n_r)$. Assume*

that there exists a multigraph H^* containing H as a spanning subgraph such that H^* is obtained by replicating some of the existing edges of H but without adding edges between nonadjacent vertices of H . Suppose furthermore that $\Delta(H^*) = \Delta(H)$ and $\Delta(G)\text{-def}(H^*) \leq n_1 \Delta(G)$. Then $\chi'_e(G) = \Delta(G)$.

Proof. Let H, H^* be as in the statement of Lemma 5. The condition

$$\Delta(G)\text{-def}(H^*) \leq n_1 \Delta(G)$$

is equivalent (arguing as in Proposition 2) to

$$|E(H^*)| \geq \frac{1}{2}(|V(H)| - n_1)\Delta(G). \quad (3)$$

By possibly removing some of the edges from H^* , we can assume that the sign of equality holds in (3) and hence that

$$|E(H^*)| = \frac{1}{2}(|V(H)| - n_1)\Delta(G), \quad (4)$$

which is equivalent to

$$\Delta(G)\text{-def}(H^*) = n_1 \Delta(G). \quad (5)$$

Let $v \in V(H) = V(H^*)$. By assumption, the edges incident with v which are in H^* but not in H are at most

$$\Delta(H) - \deg_H(v) \leq \Delta(H) - (|V(H)| - n_1) = n_1 - n_r \leq n_1 - 1,$$

so that (denoting by $\mu(H^*)$ the maximum multiplicity of the edges of H^*) we have $\mu(H^*) \leq n_1$, since H is a simple graph. But then, by Vizing's Theorem, we have

$$\chi'(H^*) \leq \Delta(H^*) + \mu(H^*) \leq \Delta(H) + n_1 = \Delta(G).$$

Thus, H^* is $\Delta(G)$ -edge colorable. But then, in particular, H^* has an equalized $\Delta(G)$ -edge coloring, which we denote by φ . It follows by (4) that each color class of φ contains exactly $\frac{1}{2}(|V(H)| - n_1)$ edges. Since $\Delta(H^*) = \Delta(H) = \Delta(G) - n_1$, it follows that every vertex of H^* misses at least n_1 of the colors given by φ .

Let ψ be the excessive coloring of H obtained by assigning to the edge $xy \in E(H)$ all the colors assigned by φ to the edges $xy \in E(H^*)$. Then clearly ψ is an excessive coloring of H using $\Delta(G)$ colors, such that each color class contains exactly $\frac{1}{2}(|V(H)| - n_1)$ edges and each vertex misses at least n_1 colors. Thus, ψ satisfies Theorem 3 and hence, by Lemma 4, Theorem 2 holds for G , as we wanted. ■

We are ready to prove the following theorem, which, together with Theorem 4, proves Theorem 2.

Theorem 5. *Let G be a 1-extendable complete multipartite graph such that $\sigma_1(G) \leq \Delta(G)$. Then $\chi'_e(G) = \Delta(G)$.*

Proof. By Lemma 5, it suffices to prove the existence of a multigraph H^* as specified in the statement of Lemma 5.

Let H^* be a maximal multigraph which is a spanning supergraph of H obtained by replicating existing edges of H without introducing edges between nonadjacent vertices of H and such that $\Delta(H^*) = \Delta(H)$. We prove that H^* satisfies the conditions of Lemma 5.

Claim 1. *There can be at most one partite set V_i of H^* containing vertices of degree less than $\Delta(H)$.*

This is obvious since otherwise we could add to H^* an edge by replicating an existent edge xy of H^* without violating the constraints on the maximum degree but contradicting the maximality of H^* .

Conclusion. If all the vertices in $V(H^*)$ have degree $\Delta(H)$ there is clearly nothing to prove, since then

$$\Delta(G)\text{-def}(H^*) = n_1|V(H)| \leq n_1\Delta(G)$$

and all the conditions of Lemma 5 are satisfied.

Thus, we can assume (by Claim 1) that there is exactly one partite set V_i of H^* containing vertices of degree less than $\Delta(H)$.

But then

$$\begin{aligned} \Delta(H)\text{-def}(H^*) &= \sum_{v \in V_i} (\Delta(H) - \deg_{H^*}(v)) \\ &\leq \sum_{v \in V_i} (\Delta(H) - \deg_H(v)) = n_i(n_i - n_r). \end{aligned}$$

Hence

$$\Delta(G)\text{-def}(H^*) \leq n_i(n_i - n_r) + n_1(|V(G)| - n_1). \tag{6}$$

Using the fact that $n_1 \geq n_i \geq n_r$, it is easily seen that

$$n_i(n_i - n_r) \leq n_1(n_1 - n_r).$$

Hence, using (6), we see that

$$\Delta(G)\text{-def}(H^*) \leq n_1(n_1 - n_r) + n_1(|V(G)| - n_1) = n_1(|V(G)| - n_r) = n_1\Delta(G).$$

Therefore, H^* satisfies all the conditions of Lemma 5 and hence Theorem 5 is proved. ■

Proof of Theorem 2. This follows immediately from Theorems 4 and 5.

ACKNOWLEDGMENT

The authors are grateful to the anonymous referees for their comments on the first version of this article.

REFERENCES

- [1] A. Bonisoli, Edge covers of K_{2n} with $2n$ one-factors, *Rendiconti del Seminario Matematico di Messina, Serie II* 9 (2003), 43–51.
- [2] A. Bonisoli and D. Cariolaro, Excessive factorizations of regular graphs, In: *Graph Theory in Paris (Proceedings of a Conference in memory of Claude Berge, Paris 2004)*, A. Bondy, J. Fonlupt, J.-L. Fouquet, J.-C. Fournier, J. L. Ramirez Alfonsin (Editors), Birkäuser, Basel, 2007, pp. 73–84.
- [3] D. G. Hoffman and C. A. Rodger, The chromatic index of complete multipartite graphs, *J Graph Theory* 16(2) (1992), 159–163.
- [4] C. J. H. McDiarmid, The solution of a timetabling problem, *J Inst Math Appl* 9 (1972), 23–34.
- [5] W. D. Wallis, *One-factorizations*, Kluwer, Dordrecht, Netherlands, 1997.
- [6] W. D. Wallis, Overfull sets of one-factors, *J Combin Math Combin Comput* 57 (2006), 151–156.
- [7] D. de Werra, Equitable colorations of graphs, *INFOR* 9 (1971), 220–237.